Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

PATIENT EDUCATION

Ethyl loflazepate

Loflazepat / Meilax / International benzodiazepine formulations

A guide for adults taking ethyl loflazepate.
Follow the directions on your prescription and your provider's recommendations.

NOT FDA-APPROVED | LONG-LASTING EFFECTS: DO NOT DRIVE

Formulations: Oral tablets or other locally approved preparations have country-specific dosing and handling instructions. This guide follows the supplied Japanese Meilax reference.

What it is used for

Ethyl loflazepate is used in Japan and some other countries for anxiety-related symptoms in specified conditions. It is not approved for medical use in the United States and is not a stand-alone treatment for major depression.

How it works

The body converts it to long-lasting metabolites that enhance GABA signaling. Effects can accumulate over days to weeks and may continue after the last dose.

What to expect over time

Effects can begin after the first dose. The timing and duration depend on the formulation and dose. Sleepiness or impaired coordination may last longer than the relief you notice.

Your provider should review whether ongoing treatment is helping and whether the dose or duration should change. Do not increase the dose when the effect seems weaker.

Taking your medication

- Follow the local prescription and do not add doses because the initial effect seems incomplete.
- If you miss a dose, take it when remembered unless the next dose is nearly due. Skip it when near the next dose; never double.
- Do not drive or operate dangerous machinery if the local instructions prohibit it.
- Consult with your provider before reducing or stopping regular use.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

ETHYL LOFLAZEPATE / 02

Side effects and safety

Possible side effects

- Sleepiness, slowed reactions, or fatigue
- Dizziness, poor balance, or falls
- Memory problems or difficulty concentrating

Keeping track of treatment

Some side effects resolve with time; others may require a change of medication. Discuss persistent or bothersome side effects with your provider.

Watch for daytime sedation, falls, memory changes, and breathing problems. Older adults and people taking other sedatives may be more sensitive.

Physical dependence can develop with regular use. Sudden stopping or a rapid reduction can cause severe withdrawal, including seizures. Dependence is different from addiction, but misuse and addiction are also possible.

If you take too much

For a suspected overdose, call Poison Control at **1-800-222-1222** immediately, even if you feel well. Call 911 for a seizure, collapse, trouble breathing, or inability to wake.

Before you start

Discuss sleep apnea, lung disease, liver or kidney disease, glaucoma, falls, and any history of substance use problems. Tell your provider about all opioids and other sedatives.

Discuss pregnancy, pregnancy plans, and breastfeeding with your provider. Treatment decisions should consider both the medication and the condition being treated.

The Japanese patient guide advises avoiding breastfeeding during treatment. Discuss a coordinated feeding and medication plan; do not abruptly stop regular use.

Medicines and interactions

Review all your medications with your provider, including over-the-counter medicines and supplements. Have new medications checked before taking them.

Combining a benzodiazepine with an opioid can cause profound sedation, slowed breathing, coma, or death. Use such combinations only under a specifically reviewed treatment plan.

Avoid alcohol. Sleeping pills, sedating antihistamines, muscle relaxants, and other sedatives can increase impairment and breathing risks.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

ETHYL LOFLAZEPATE / 03

Living with your treatment

When to get help

Emergency symptoms. Call 911 for slow or difficult breathing, inability to wake, collapse, or a seizure. Severe confusion or hallucinations after reducing or stopping may be dangerous withdrawal.

Prompt medical advice. Contact your provider promptly for repeated falls, marked memory problems, unusual aggression, or needing more medication to get the same effect.

If you feel unsafe or have thoughts of self-harm, call 911 or go to an emergency department.

Your questions answered

Can I stop after taking it regularly?
Consult with your provider before reducing or stopping. A gradual, individualized taper helps limit withdrawal. Do not borrow medication or change the schedule to manage withdrawal yourself.

What if I feel unusually agitated?
Some people develop disinhibition, irritability, or agitation. Contact your provider rather than taking an extra dose.

Everyday tips

Do not drive until you know how the medication affects you. Avoid driving whenever sleepy or impaired.

Keep medication locked away. Never share it, even with someone who has similar symptoms.

Arrange refills or taper instructions before you run out.

Storage and safety

Keep medication in its labeled container, away from excess heat and moisture and out of reach of children and pets. Do not share it. Ask a pharmacy about disposal of unused medication.

Sources

PMDA: Meilax patient guide, 2023

Educational handout; does not replace your prescription or provider's recommendations.