Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

PATIENT EDUCATION

Mianserin

International antidepressant / UK labeling reference

A guide for adults taking mianserin.
Follow the directions on your prescription and your provider's recommendations.

KEEP BLOOD-TEST APPOINTMENTS

FEVER, SORE THROAT, OR MOUTH ULCERS: ACT PROMPTLY

Formulations: Tablets may be taken at bedtime or in divided doses as prescribed. Blood-count monitoring is part of the treatment plan.

What it is used for

Mianserin treats depressive illness in countries where it is supplied. It is not FDA-approved. This guide follows the reviewed UK oral labeling.

How it works

It affects norepinephrine and serotonin signaling by blocking several receptors. Histamine-receptor effects contribute to drowsiness and increased appetite.

What to expect over time

Sleep, energy, or appetite may change before mood improves. Benefit usually builds over several weeks; some conditions need a longer trial at a useful dose.

Track the symptoms you are treating and how you manage daily activities. Keep follow-up appointments. Schedule an earlier appointment if your mood gets worse.

Taking your medication

- Swallow tablets whole with water.
- Avoid alcohol and follow the leaflet's driving restrictions.
- If you miss a dose, take it when you remember unless it is almost time for the next dose. Then skip it. Do not double a dose.
- Consult with your provider before routine stopping. Infection symptoms require immediate action.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

MIANSERIN / 02

Side effects and safety

Possible side effects

- Drowsiness, dizziness, or low blood pressure
- Dry mouth or constipation
- Increased appetite or weight

Keeping track of treatment

Some side effects resolve with time; others may require a change of medication. Discuss persistent or bothersome side effects with your provider.

Antidepressants can increase suicidal thoughts in people under 25. Watch for changes in mood or behavior at any age, especially when starting treatment or changing a dose.

Tell your provider about emotional numbness, unusual agitation, or a reduced need for sleep with increased energy. These changes may call for a treatment adjustment.

UK labeling recommends a full blood count every four weeks during the first three months. Infection symptoms still need prompt assessment between tests and afterward.

If you take too much

For a suspected overdose, call Poison Control at **1-800-222-1222** immediately, even if you feel well. Call 911 for a seizure, collapse, trouble breathing, or inability to wake.

Before you start

Discuss any history of mania or bipolar disorder, seizures, heart disease, glaucoma, and kidney or liver problems.

Discuss pregnancy, pregnancy plans, and breastfeeding with your provider. Treatment decisions should consider both the medication and the condition being treated.

Severe liver disease and mania rule out use under the reviewed UK label. Discuss prior blood disorders and unexplained infections.

Medicines and interactions

Review all your medications with your provider, including over-the-counter medicines and supplements. Have new medications checked before taking them.

MAOI antidepressants require a medication-specific waiting period before or after this treatment. Linezolid and IV methylene blue also need special precautions. Consult with your provider before any switch.

Other medicines that increase serotonin can raise the risk of serotonin syndrome. Alcohol and sedating medicines may worsen dizziness or impairment.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

MIANSERIN / 03

Living with your treatment

When to get help

Emergency symptoms. Get emergency care for a seizure, severe allergic reaction, or sudden eye pain with vision changes. Fever with agitation, diarrhea, and stiff or twitching muscles may be serotonin syndrome and needs urgent assessment.

Prompt medical advice. Stop mianserin and seek urgent medical advice for fever, sore throat, mouth ulcers, or other infection signs; a blood count is needed. Report jaundice, unusual bruising, or new manic symptoms promptly.

If you feel unsafe or have thoughts of self-harm, call 911 or go to an emergency department.

Your questions answered

Can I stop when I feel better?
Consult with your provider. Continuing treatment after improvement can reduce the chance that symptoms return. Most antidepressants should be reduced gradually to limit withdrawal symptoms.

What if I have sexual side effects?
Changes in desire, arousal, or orgasm are worth discussing. Your provider can review the dose or treatment options; do not stop or skip doses on your own.

Everyday tips

A regular dose time and refill reminder can help prevent missed doses.

Bring a few specific observations to appointments: sleep, motivation, anxiety, and activities that have become easier or harder.

Storage and safety

Keep medication in its labeled container, away from excess heat and moisture and out of reach of children and pets. Do not share it. Ask a pharmacy about disposal of unused medication.

Sources

Official prescribing information: mianserin

Educational handout; does not replace your prescription or provider's recommendations.