Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

PATIENT EDUCATION

Zotepine

Lodopin / International antipsychotic formulations

A guide for adults taking zotepine.
Follow the directions on your prescription and your provider's recommendations.

FOLLOW YOUR EXACT DOSE SCHEDULE

REPORT SEVERE CONSTIPATION

Formulations: Oral tablets and granules differ in strength and measurement. Follow the prescribed divided-dose schedule and locally supplied instructions.

What it is used for

Zotepine is used in some countries for schizophrenia. It is not FDA-approved. This guide follows Japanese Lodopin information; use the locally supplied prescription and leaflet for dosing.

How it works

It affects dopamine and serotonin receptors and norepinephrine signaling. Effects on other receptors can cause sedation, constipation, and weight changes.

What to expect over time

Some symptoms improve earlier than others. Changes in hallucinations, mood, or organized thinking often develop over several weeks. Track practical goals, such as regular sleep and daily activities, and report side effects or limited benefit.

Taking your medication

- Take only the prescribed amount and schedule. Ask a pharmacist to demonstrate granule measurement.
- Do not substitute tablets and granules by appearance or spoonful.
- If you miss a dose, take it when you remember unless it is almost time for the next dose. Then skip it. Do not double a dose.
- Follow local restrictions on driving, work at heights, and dangerous machinery.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

ZOTEPINE / 02

Side effects and safety

Possible side effects

- Sleepiness or dizziness
- Dry mouth, constipation, or blurred vision
- Weight gain
- Restlessness, stiffness, or tremor

Keeping track of treatment

Some side effects resolve with time; others may require a change of medication. Discuss persistent or bothersome side effects with your provider.

Expect periodic checks of weight, blood pressure, blood sugar, cholesterol, and movement symptoms. Some changes, such as higher blood sugar, may occur before you feel different.

Report new mouth, tongue, face, or limb movements. Tardive dyskinesia is a movement disorder that can persist after medication is stopped.

Track bowel function closely; severe constipation can become dangerous. Weight, glucose, lipids, ECG, blood counts, and liver tests may be monitored.

If you take too much

For a suspected overdose, call Poison Control at **1-800-222-1222** immediately, even if you feel well. Call 911 for a seizure, collapse, trouble breathing, or inability to wake.

Before you start

Discuss diabetes, heart-rhythm problems, fainting, seizures, low blood counts, swallowing difficulties, and kidney or liver disease. Antipsychotics increase the risk of death in older adults with dementia-related psychosis.

Discuss pregnancy, pregnancy plans, and breastfeeding with your provider. Treatment decisions should consider both the medication and the condition being treated.

Discuss seizure history, severe constipation, urinary obstruction, heart disease, gout, kidney stones, and liver or kidney impairment.

Medicines and interactions

Review all your medications with your provider, including over-the-counter medicines and supplements. Have new medications checked before taking them.

Alcohol, opioids, sleeping pills, and sedating allergy medicines can increase impairment. Blood-pressure medicines may add lightheadedness or falls.

Medicines that affect heart rhythm or dopamine signaling need particular review. Some antibiotics, antifungals, and seizure medicines change antipsychotic levels.

Jeffrey Panko NP
PSYCHIATRY

Jeffrey Panko, PMHNP-BC
Phone 646-549-4694 | Fax 646-558-4238
JMPankoNP@JeffreyPanko.com
jeffreypanko.com

ZOTEPINE / 03

Living with your treatment

When to get help

Emergency symptoms. Get emergency care for fever with severe muscle stiffness and confusion, a seizure, fainting with an irregular heartbeat, or trouble breathing. Fever and stiffness can signal neuroleptic malignant syndrome, a rare but serious reaction.

Prompt medical advice. Seek urgent care for severe constipation with abdominal pain or swelling, inability to pass gas, fever, new movements, or urinary retention.

If you feel unsafe or have thoughts of self-harm, call 911 or go to an emergency department.

Your questions answered

Can I stop when I feel better?
Consult with your provider. Continuing treatment may prevent symptoms from returning. Abrupt stopping can cause withdrawal symptoms or a relapse.

What if I cannot sit still?
An inner urge to keep moving can be akathisia, a medication side effect. Tell your provider promptly; it can be mistaken for worsening anxiety.

Everyday tips

Stand up slowly. Avoid driving until you know how the medication affects alertness and coordination.

Stay hydrated and take breaks in hot weather. Some antipsychotics interfere with temperature regulation.

Discuss appetite or weight changes early, and keep scheduled monitoring visits.

Storage and safety

Keep medication in its labeled container, away from excess heat and moisture and out of reach of children and pets. Do not share it. Ask a pharmacy about disposal of unused medication.

Sources

PMDA: Lodopin patient guide

Educational handout; does not replace your prescription or provider's recommendations.